

PIN OAK

***Quercus palustris* Muenchh.**

plant symbol = QUPA2

Contributed by: USDA NRCS Northeast Plant Materials Program



Robert H. Mohlenbrock
 USDA NRCS 1991. Southern Wetland Flora
 @USDA NRCS PLANTS

Uses

Even though the wood is heavy, strong, and hard, pin oak is typically used for fuel wood, wood pulp, and railroad ties, since it tends to warp. Due to this species' form, adaptability, growth rate, longevity, and fall foliar display, it is popular for ornamental usage. Pin oak is utilized by many game species, especially wood ducks, white-tail deer, and wild turkey. Due to its use by these highly sought after animals, pin oak is commonly planted for food plots.

Status

Please consult the PLANTS Web site and your State Department of Natural Resources for this plant's current status (e.g. threatened or endangered species, state noxious status, and wetland indicator values).

Description

Pin oak is a moderately large tree with normal heights ranging from 70 to 90 feet with diameters between 2 and 3 feet. Trees reaching 120 feet tall with 5-foot diameters are occasionally encountered on good sites. The bark of this tree is smooth, reddish to grayish-brown during the juvenile period, becoming darker and shallowly fissured as the tree growth slows with age. The lower branches of pin oak are prostrate to

descending, with smooth, slender, reddish-brown twigs. Clusters of pointed buds are located at the tips of twigs. Three to five inch alternate leaves have 5 to 7 points or lobes with bristled tips and deep C-shaped sinuses. The leaves change in color from a dark green to a deep scarlet red in fall. The leaves are deciduous but will usually persist on the tree into winter.


The flowers of pin oak emerge soon after new leaves unfold in spring (April to mid-May). The acorns that develop are roundish, short stalked, 3/8 to 1/2 inches long, and capped with a thin and shallow saucer-like cup. The acorns will take 16 to 18 months to develop from pollination to maturity. When mature the acorn turns light brown to reddish-brown, and will drop from September to November. In 30 to 35 year old stands of pin oak, 4,000 to 20,000 sound acorns per acre yields have been documented. There are 410 acorns per pound.

Pin oak is often confused with scarlet oak (*Q. coccinea*) due to similar appearance. Scarlet oak is an upland species that prefers soils with good drainage on dry sites.

Adaptation

Pin oak's native range spans from Rhode Island, Pennsylvania, Michigan, to Kansas south to North Carolina and northern Arkansas. This tree grows under a wide range of site conditions, but is a true bottomland tree. It is seldom found growing above elevations of 800 feet, or on sloped ground. Pin oak grows in practically pure stands with good growth rates on wet, shallow sites with heavy soils that drain poorly. On better quality sites it will often grow larger, but is normally outcompeted by other species.

Known Distribution



01/08/01

Establishment

Both seed and stump sprouts are sources of natural regeneration for pin oak. Although viable acorns sink, this species is often dispersed by water, as well as animals, wind, and gravity. In areas that are regularly flooded, acorn damage by insects is reduced and germination is typically faster than most other species. Since there is more than adequate annual seed yields and adequate moisture on the forest floor in spring, it is not uncommon to see seedlings blanketing the ground under pin oak.

Under nursery production conditions, mature acorns are placed on raised beds in fall or spring, but germination will not occur until spring. Shoot elongation begins about the same time as leaf out of established trees and continues until fall. Seedlings are distributed by most nurseries as 1 year old bareroot stock for field plantings, but balled and burlapped saplings are distributed for ornamental uses.

Management

This section is under development.

Pests and Potential Problems

Insect and disease damage to pin oak is not common, but entire stands may be killed by prolonged flooding.

Cultivars, Improved, and Selected Materials (and area of origin)

‘Crownright,’ ‘Sovereign.’ The species is available through numerous native plant nurseries. Nursery produced seedlings are currently grown from local or regional seed sources.

Prepared By & Species Coordinator:

USDA NRCS Northeast Plant Materials Program

12Jan2001 JLK

For more information about this and other plants, please contact your local NRCS field office or Conservation District, and visit the PLANTS <<http://plants.usda.gov>> and Plant Materials Program Web sites <<http://Plant-Materials.nrcs.usda.gov>>.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call (202) 720-5964 (voice or TDD). USDA is an equal opportunity provider and employer.